

AUTOMATIC DETECTION OF VOLCANIC THERMAL FEATURES USING SATELLITE OBSERVATIONS (V35E-0176)

Corradino, C.^{1,2}, Ramsey, M.S.², Leggett, T.N.², Del Negro, C.¹

¹Istituto Nazionale di Geofisica e Vulcanologia (INGV), Catania, Italy, ²Department of Geology and Environmental Science, University of Pittsburgh, Pittsburgh, PA, USA

AGU FALL MEETING - New Orleans, LA, 13-17 December 2021

Session: V029 - Volcano Hazard Monitoring from Space using Statistical Methods and Machine Learning

MOTIVATIONS

Satellite observations are widely used to investigate, monitor, and forecast volcanic activity. Spaceborne thermal infrared (TIR) measurements of thermal volcanic features improve our understanding of the underlying processes leading to eruptions[1-3]. Unfortunately, although different approaches have been proposed to detect and estimate the temperature above background of volcanic thermal anomalies, detecting subtle changes is still a challenging task. Here, we develop an artificial intelligence (AI) approach to automatically detect volcanic thermal features by using both low-spatial, high-temporal resolution MODerate-resolution Imaging Spectroradiometer (MODIS) data to detect hotter thermal features at short time scales; as well as high-spatial, low-temporal resolution Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER) to detect more subtle thermal changes otherwise missed by MODIS.

SPACEBORNE THERMAL INFRARED SENSORS

HIGH SPATIAL RESOLUTION: ASTER

ASTER is on board of Terra satellite. It has a revisit time of 16 days at the equator, but with the URP program started in 2009, it acquires data much more rapidly.

- TIR bands [8.125-11.65] at 90m spatial resolution are available

- Topographically corrected images AST_L1T are used




Fig.1. ASTER acquisition over Lascar, 21/10/05 03:16 (a) and Etna - 06/09/04 21:02 (b)

HIGH TEMPORAL RESOLUTION: MODIS

MODIS is on board of Terra and Aqua satellites. The MODIS sensors are viewing the entire Earth's surface up to 4 times per day.

- Images acquired at 36 spectral bands at 1km are available

- Level 2 LST (Land Surface Temperature) is used




Fig. 2. MODIS acquisition over Lascar - 21/10/05 03:16 (a) and Etna - 06/09/04 21:02 (b)

ARTIFICIAL INTELLIGENCE TECHNIQUES FOR VOLCANIC THERMAL FEATURE DETECTION

SPATIAL VOLCANIC THERMAL FEATURE EXTRACTION USING DEEP LEARNING (DL)

Supervised techniques require a large volume of training labeled data. Here we use a statistical approach (more details in V35E-0178) to label data followed by an expert supervision step required to check correctly labeled images, which serve as a target. Input and target data are then processed and fed into a UNET that will learn volcanic thermal features against background, clouds and missing data (i.e. border pixels with missing values).




Fig. 3. Workflow of the proposed deep learning (DL) approach to retrieve volcanic spatial thermal features.

TEMPORAL VOLCANIC THERMAL FEATURE EXTRACTION USING ARTIFICIAL INTELLIGENCE (AI)

- Select volcano coordinate (Lat, Lon)

- Select a buffer around volcano coordinate
- Select date interval
- Apply a cloud coverage filter
- LST conversion from Kelvin to Celsius

- A simple regression model (OLS) is applied to fit LST data time series 'normal' trend for each pixel
- The reconstruction error is computed as the deviation of the real LST time series from the normal reconstructed trend

- High reconstruction errors are related to abnormal thermal behaviours
- An isolation random forest is used as unsupervised machine learning (ML) approach to detect anomalies based on the reconstruction errors




Fig. 4. Workflow of the proposed artificial intelligence (AI) approach to retrieve temporal volcanic thermal features. Graphs show the time series of the maximum values inside the region of interest


RESULTS

TEMPORAL VOLCANIC THERMAL FEATURES EXTRACTION USING AI

MODIS: Etna (2000 to 2021)




MODIS: Kiliuevskoi (2000 to 2021)




SPATIAL VOLCANIC THERMAL FEATURES EXTRACTION USING DL


ASTER: Etna - 2013/05/18




ASTER: Fuego - 2003/10/14




ASTER: Kiliuevskoi - 2018/01/11





ASTER: Lascar - 2017/09/01



ASTER: Popocatepetl - 2017/07/26



- ASTER and MODIS data are shown together
- For each month, the maximum value of the reconstruction error (a) and the detected anomalies (b) are shown
- For some eruptions (blue arrows), a thermal increase occurs before an eruption (gray shaded area)



CONCLUSIONS

An AI approach has been designed to automatically extract volcanic features both spatially (exploiting high spatial resolution satellite thermal data using convolutional neural networks) and temporally (exploiting high temporal resolution satellite data using the isolation random forest as anomaly detection algorithm). With these techniques, very large datasets can be easily processed and high temporal and high spatial resolution satellite data combined to improve thermal volcanic precursory monitoring by detecting very low-level anomalies linked to pre-eruptive activity.

REFERENCES

1. Murphy, S. W., et al. "MODIS and ASTER synergy for characterizing thermal volcanic activity." Remote sensing of environment 131 (2013): 195-205.
2. Reath, Kevin, et al. "Thermal, deformation, and degassing remote sensing time series (CE 2000–2017) at the 47 most active volcanoes in Latin America: implications for volcanic systems." Journal of Geophysical Research: Solid Earth 124.1 (2019): 195-218.
3. Girona, T., Realmuto, V. & Lundgren, P. Large-scale thermal unrest of volcanoes for years prior to eruption. Nat. Geosci. 14, 238–241 (2021).

AKNOWLEDGEMENTS

This research was partially funded by the Fulbright Visiting Research Program awarded to CC and the NASA Science of Terra, Aqua, and Suomi NPP Program (80NSSC18K1001) awarded to MSR.